


Childhood Immunization Checklist



The U.S. Department of Health and Human Services' Centers for Disease Control and Prevention recommend that children age 18 and younger at a normal level of health risk receive the following immunizations. Some immunization regimens may need to be modified, postponed, or omitted entirely based on the patient's health risk factors. Each parent should speak to their child's pediatrician or other primary care provider.

- Diphtheria, tetanus, and pertussis (DTaP)**
Five doses, with one dose each at 2 months, 4 months, 6 months, 15 – 18 months, and 4 – 6 years
- Haemophilus influenzae type B**
Four doses, with one dose each at 2 months, 4 months, 6 months (as needed depending on the brand of vaccine), and 12 – 15 months
- Hepatitis A**
Two doses, with one dose each at 12 – 23 months, second dose at least six months later
- Hepatitis B**
Three doses, with one dose each at birth, 1 – 2 months, and 6 – 18 months
- Human papillomavirus**
If 9 – 14 years old at first vaccination:
Two doses, administered six to 12 months apart
If 15 or older at first vaccination:
Three doses, with the second dose one to two months after the first and the third dose six months following the second dose
- Inactivated poliovirus**
Four doses, with one dose each at 2 months, 4 months, 6 – 18 months, and 4 – 6 years
- Influenza**
One dose annually by the end of October, beginning at age 6 months
- Measles, mumps, and rubella**
Two doses, with one dose each at 12 – 15 months and 4 – 6 years (infants ages 6 – 11 months should have one dose before traveling abroad)
- Meningococcal serogroup A, C, W, Y**
Two doses, with one dose each at 11 – 12 years and 16 years
- Meningococcal serogroup B**
Two doses at 16 – 18 years
- Pneumonia**
Four doses, with one dose each at 2 months, 4 months, 6 months, and 12 – 15 months
- Rotavirus**
Two doses, with one dose each at 2 months and 4 months (Rotarix®), or three doses, with one dose each at 2 months, 4 months, and 6 months (RotaTeq®)
- Tetanus, diphtheria, and pertussis (Tdap)**
One dose at 11 – 12 years
- Varicella**
Two doses, with one dose each at 12 – 15 months and 4 – 6 years

